Homework 5: Linear Independence

Instructions. In Problems 2 and 3, make sure to use the definition of linear independence given in class. It is different from the textbook's definition. Assignments should be **stapled** and written clearly and legibly.

1. $\S4.3, \#3(a), 4(a), 15(b), 16(a)(c)$.

Try to solve # 15(b) and 16(a) with as few computations as possible, and try to be fairly rigorous for 16(c).

2. Let $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4\}$ be a linearly independent set of vectors in a vector space V. Using only the definition of linear independence, prove that $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ is linearly independent as well.

Hint. Begin the proof as follows:

"Suppose that $k_1\mathbf{v}_1 + k_2\mathbf{v}_2 + k_3\mathbf{v}_3 = \mathbf{0}$ for some scalars k_1, k_2, k_3 . I must show that $k_1 = k_2 = k_3 = 0$."

3. Using only the definition of linear independence, prove that if $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ is linearly independent, then so is $\{\mathbf{u} + \mathbf{v}, \mathbf{u} + 2\mathbf{w}, \mathbf{v} + 3\mathbf{w}\}$.

Hint. Begin the proof as follows:

"Suppose that $c(\mathbf{u} + \mathbf{v}) + d(\mathbf{u} + 2\mathbf{w}) + e(\mathbf{v} + 3\mathbf{w}) = \mathbf{0}$ for some scalars c, d, e. I must show that c = d = e = 0."